

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE (DD-MM-YYYY) 16-03-2006		2. REPORT TYPE Journal Article		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE  Preparation , Characterization, and Crystal Structures of the SO <sub>3</sub> NHF <sup>-</sup> and SO <sub>3</sub> NF <sub>2</sub> <sup>-</sup> Anions (POSTPRINT)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Ralf Haiges, Ross Wagner, Muhammed Yousufuddin, Markus Etzkorn, G.K. Surya Prakash, & Karl O. Christe (USC); Jerry A. Boatz (AFRL/PRSP); Robert D. Chapman (NAWC); Mark F. Welker & Charles B. Kreutzberger (TPL, Inc.)				5d. PROJECT NUMBER	
				5e. TASK NUMBER 23030423	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/PRSP 10 E. Saturn Blvd. Edwards AFB CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NUMBER  AFRL-PR-ED-JA-2006-101	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-PR-ED-JA-2006-101	
12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited (AFRL-ERS-PAS-2006-084).					
13. SUPPLEMENTARY NOTES ©2006Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim. Published in Angewandte Chemie Int. Ed. 2006, 45, 5179-5184.					
14. ABSTRACT Recently, a new class of high-energy-containing materials, <i>gem</i> -bis(difluoramino)-substituted heterocyclic nitramines, has gained attention as high-energy oxidizers: HNF <sup>1,2</sup> and TNF <sup>3</sup> have been successfully synthesized under strongly acidic conditions from their corresponding ketone derivatives using an excess of difluoramine. <sup>4</sup> HNF <sub>2</sub> is an unpredictably shock-sensitive and thermally unstable, gaseous compound <sup>5,6</sup> which can be generated from different precursors, e.g., tetrafluorohydrazine, <sup>7</sup> <i>N,N</i> -difluorourea, <sup>8</sup> <i>N,N</i> -difluorocarbamates, <sup>9</sup> or trityldifluoramine. <sup>10</sup> Out of these precursors, only trityldifluoramine is a stable storable solid. However, it is not useful as a general reagent for the preparation of larger quantities of <i>gem</i> -bis(difluoramines) because its synthesis requires the use of expensive N <sub>2</sub> F <sub>4</sub> which is commercially unavailable and must be prepared from difluoramine, and of equivalent amounts of mercury in an organic solvent. The use of mercury presents environmental problems, and working with N <sub>2</sub> F <sub>4</sub> in an organic solvent can be hazardous. Therefore, it is highly desirable to develop a stable, solid, readily accessible difluoramine source. Obvious candidates for HNF <sub>2</sub> sources were difluorosulfamate salts. Although the parent free acid, HOSO <sub>2</sub> NF <sub>2</sub> , had been known since 1961 and has been widely used as a difluoroaminating reagent, <sup>11,12</sup> no reports could be found on the existence of its salts. In this paper, we report the results from two independent studies.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  SAR	18. NUMBER OF PAGES  7	19a. NAME OF RESPONSIBLE PERSON Dr. Scott A. Shackelford
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)
Unclassified	Unclassified	Unclassified			N/A

DOI: 10.1002/anie.200601020

# Preparation, Characterization, and Crystal Structures of the $\text{SO}_3\text{NHF}^-$ and $\text{SO}_3\text{NF}_2^-$ Ions\*\*

Ralf Haiges,\* Ross Wagner, Jerry A. Boatz,  
Muhammed Yousufuddin, Markus Etzkorn,  
G. K. Surya Prakash, Karl O. Christe,\*  
Robert D. Chapman,\* Mark F. Welker, and  
Charles B. Kreutzberger

*Dedicated to Dr. Carl Schack  
on the occasion of his 70th birthday*

Recently, a new class of materials having high-energy, *gem*-bis(difluoroamino)-substituted heterocyclic nitroamines, has gained attention for use as high-energy oxidizers: HNF<sup>+</sup><sub>2</sub><sup>[1,2]</sup> and TNF<sup>+</sup><sub>2</sub><sup>[3]</sup> have been synthesized under strongly acidic conditions from their corresponding ketone derivatives by using an excess of difluoroamine.<sup>[4]</sup>

HNF<sub>2</sub> is a shock-sensitive and thermally unstable gaseous compound,<sup>[5,6]</sup> which can be generated from different precursors, for example, tetrafluorohydrazine,<sup>[7]</sup> *N,N*-difluoro-urea,<sup>[8]</sup> *N,N*-difluorocarbamates,<sup>[9]</sup> or trityldifluoroamine.<sup>[10]</sup> Of these precursors, only trityldifluoroamine is a stable,

[\*] Dr. R. Haiges, Dr. R. Wagner, M. Yousufuddin, Dr. M. Etzkorn,  
Prof. Dr. G. K. S. Prakash, Prof. Dr. K. O. Christe  
Loker Research Institute and Department of Chemistry  
University of Southern California  
Los Angeles, CA 90089-1661 (USA)  
Fax: (+1) 213-740-6679  
E-mail: haiges@usc.edu  
kchriste@usc.edu

Dr. R. D. Chapman  
Chemistry Branch (Code 498200D)  
Naval Air Warfare Center Weapons Division  
1900 N. Knox Rd. Stop 6303  
China Lake, CA 93555-6106 (USA)  
Fax: (+1) 760-939-1617  
E-mail: robert.chapman@navy.mil


Dr. J. A. Boatz  
Space and Missile Propulsion Division  
Air Force Research Laboratory (AFRL/PRSP)  
10 East Saturn Boulevard, Bldg 8451  
Edwards Air Force Base, CA 93524 (USA)

Dr. M. F. Welker, Dr. C. B. Kreutzberger  
TPL, Inc.  
3921 Academy Parkway North NE  
Albuquerque, NM 87109-4416 (USA)

[\*\*] The work at the University of Southern California (USC) was funded by the Air Force Office of Scientific Research, the Office of Naval Research, Lawrence Livermore National Laboratory, and the National Science Foundation, and that at the Naval Air Warfare Center Weapons Division (NAWCWD) and TPL, Inc. was supported by the Office of Naval Research. Portions of this work were previously reported in limited-distribution presentations.<sup>[1]</sup>




Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.



storable solid. However, it is not useful as a general reagent for the preparation of larger quantities of *gem*-bis(difluoroamines) because its synthesis requires the use of expensive  $\text{N}_2\text{F}_4$ , which is not commercially available and must be prepared from difluoroamine, and of equivalent amounts of mercury in an organic solvent. The use of mercury presents environmental problems, and working with  $\text{N}_2\text{F}_4$  in an organic solvent can be hazardous. Therefore, it is highly desirable to develop a stable, solid, readily accessible difluoroamine source. Obvious candidates for  $\text{HNF}_2$  sources are difluoro-sulfamate salts. Although the parent free acid  $\text{HOSO}_2\text{NF}_2$  had been known since 1961 and has been widely used as a difluoroaminating reagent,<sup>[11,12]</sup> no reports could be found on the existence of its salts. Herein, we report the results from two independent studies.

$\text{Na}^+\text{SO}_3\text{NF}_2^-$  was isolated by fluorinating aqueous sulfamic acid at  $0^\circ\text{C}$ , neutralizing the product with concentrated aqueous sodium hydroxide, filtering off the less-soluble sodium fluoride by-product, and drying the neutral solution under vacuum. This process was improved by direct fluorination of  $\text{SO}_3\text{NH}_2^-$  salts with diluted elemental fluorine in aqueous solutions at  $0^\circ\text{C}$  [Eq. (1)].



The reaction conditions were similar to those previously used for the fluorination of urea,<sup>[8a]</sup> carbamates,<sup>[9a]</sup> and sulfamide.<sup>[13]</sup> The acidic reaction mixture was not stable above approximately  $5^\circ\text{C}$ , and the difluorosulfamate [Eq. (2)] rapidly hydrolyzed upon warming to ambient temperature.



The ease of this acid-catalyzed hydrolysis can account for the lack of previous reports of isolated salts of the difluoro-sulfamate anion. The isolation of difluorosulfamate salts required careful control of the reaction conditions and rapid removal of the water and HF by-product at  $0^\circ\text{C}$  in a vacuum. This method resulted in the isolation of pure, colorless  $\text{NaSO}_3\text{NF}_2$  in 94% yield.

The dry sodium salt is stable at room temperature but decomposes when exposed to atmospheric moisture. It was stored in the dry argon atmosphere of a glove box for a period of 4 months without showing any sign of decomposition. The identity of the compound was established by the observed material balance, vibrational and multinuclear NMR spectroscopy, electronic structure calculations, and by converting

it into  $[\text{PNP}][\text{SO}_3\text{NF}_2] \cdot \text{CH}_2\text{Cl}_2$  (PNP = bis(triphenylphosphoranylidene)ammonium) and determining its crystal structure.

The Raman and IR spectra of solid  $\text{NaSO}_3\text{NF}_2$  are shown in the Supporting Information, and the observed frequencies and intensities are listed in Table 1. They were assigned by

**Table 1:** Comparison of observed and unscaled calculated MP2/6-311 + G(d) vibrational frequencies ( $\text{cm}^{-1}$ ) and intensities for  $\text{SO}_3\text{NF}_2^-$  in point group  $C_s$ .

Mode	Approx. mode description	Observed <sup>[a], [e]</sup>		Calculated <sup>[b]</sup>
		IR	Raman	
a'	$\nu_1 \nu_{\text{as}} \text{SO}_3$	1309 vs	1304 [0.8]	1286 (364) [7.2 dp]
	$\nu_2 \nu_{\text{s}} \text{SO}_3$	1080 s	1089 [10.0]	1051 (59) [38 p]
	$\nu_3 \nu_{\text{s}} \text{NF}_2$	968 w	997 [4.0] 971 [0.9]	979 (72) [4.4 p]
	$\nu_4 \nu \text{SN}$	715 m	722 [5.5]	671 (64) [17 p]
	$\nu_5 \delta_{\text{as}} \text{SO}_3$	627 s 617 s	630 [1.1]	601 (140) [0.6 dp]
	$\nu_6 \delta_{\text{sciss}} \text{NF}_2$		526 [2.6]	521 (1.6) [7.3 p]
	$\nu_7 \delta_{\text{sym}} \text{SO}_3$	525 vw		501 (25) [4.6 p]
	$\nu_8 \delta_{\text{rock}} \text{SO}_3$	[c]	337 [4.3]	298 (1.6) [8.6 p]
	$\nu_9 \delta_{\text{wag}} \text{NF}_2$	[c]	264 [1.3]	224 (0.5) [2.4 p]
	$\nu_{10} \nu_{\text{as}} \text{SO}_3$	1284 vs	1267 [0.7]	1282 (362) [7.7 dp]
a''	$\nu_{11} \nu_{\text{as}} \text{NF}_2$	869 m	868 [1.6]	829 (75) [2.1 dp]
	$\nu_{12} \delta_{\text{as}} \text{SO}_3$	532 w	537 [2.0]	532 (35) [3.5 dp]
	$\nu_{13} \delta_{\text{rock}} \text{SO}_3$	[c]	390 [2.6]	325 (0.8) [0.8 dp]
	$\nu_{14} \delta_{\text{twist}} \text{NF}_2$	[c]	186 [1.3]	154 (0.02) [6.1 dp]
	$\tau \text{S-N}$	[c]	[d]	37 (0.00) [2.0 dp]

[a] As  $\text{Na}^+$  salt. Relative Raman intensities given in brackets. [b] IR intensities given in  $\text{kmol}^{-1}$  (in parentheses) and Raman intensities given in  $\text{\AA}^4 \text{amu}^{-1}$  (in brackets). [c] Not observed, IR spectrum recorded only between 4000 and  $400 \text{ cm}^{-1}$ . [d] Not observed, Raman spectrum recorded only between 3600 and  $80 \text{ cm}^{-1}$ . [e] In addition to the listed bands, IR bands at 1627 m, 1209 w, 1184 m, 565 w, and Raman bands at 1618 [0.5], 1321 [0.7], 572 [1.2]  $\text{cm}^{-1}$  were observed but were not assigned.


comparison with those calculated at the MP2<sup>[14]</sup> level of theory with the 6-311 + G(d) basis set. Whereas the  $^{19}\text{F}$  NMR spectrum of naturally abundant  $[\text{N}^{14}]\text{NaSO}_3\text{NF}_2$  in  $\text{CD}_3\text{CN}$  gives only a broad resonance at  $\delta = 34 \text{ ppm}$ , the spectrum of an  $^{15}\text{N}$ -labeled sample in the same solvent shows a sharp doublet at  $\delta = 33.8 \text{ ppm}$  with  $^1J(^{19}\text{F}-^{15}\text{N}) = 138 \text{ Hz}$ . The  $^{15}\text{N}$  NMR spectrum shows a sharp triplet at  $\delta = -20.4 \text{ ppm}$  with the same  $^1J(^{15}\text{N}-^{19}\text{F})$  coupling constant.

Colorless and air-stable  $[\text{PNP}][\text{SO}_3\text{NF}_2]$  was obtained by neutralizing the fluorination reaction mixture, adding  $\text{PNP}^+\text{Cl}^-$ , and extracting with methylene chloride [Eq. (3)]. Single crystals of  $[\text{PNP}][\text{SO}_3\text{NF}_2] \cdot \text{CH}_2\text{Cl}_2$  suitable for X-ray crystal structure determination were obtained by recrystallization from  $\text{CH}_2\text{Cl}_2$ .



$[\text{PNP}][\text{SO}_3\text{NF}_2] \cdot \text{CH}_2\text{Cl}_2$  crystallizes in the triclinic space group  $P\bar{1}$ . The X-ray structure analysis reveals the presence of two isolated  $[\text{PNP}]^+$  and  $[\text{SO}_3\text{NF}_2]^-$  units together with two disordered  $\text{CH}_2\text{Cl}_2$  molecules in the unit cell (packing diagrams are given in the Supporting Information). The

closest F...H and O...H contacts between neighboring cations and anions are 2.682 Å and 2.349 Å, respectively. The dimensions of the SO<sub>3</sub>N skeleton of the SO<sub>3</sub>NF<sub>2</sub><sup>−</sup> ion (Figure 1) are more similar to those of sulfamic acid<sup>[15]</sup> than




**Figure 1.** ORTEP drawing of the anion of [PNP][SO<sub>3</sub>NF<sub>2</sub>]·CH<sub>2</sub>Cl<sub>2</sub>. Thermal ellipsoids are shown at the 50% probability level. Bond lengths [Å] and angles [°]: S–O1 1.428(4), S–O2 1.424(3), S–O3 1.424(3), S–N 1.772(4), F1–N 1.412(5), F2–N 1.463(6); O1–S–O2 114.7(2), O1–S–O3 116.1(2), O2–S–O3 116.8(2), O1–S–N 100.1(2), O2–S–N 106.1(2), O3–S–N 99.3(2), F1–N–F2 99.0(3), F1–N–S 104.5(3), F2–N–S 99.6(3).

to the ones reported for salts with the SO<sub>3</sub>NH<sub>2</sub><sup>−</sup> ion.<sup>[16]</sup> Estimations of the double-bond character of the S–N bond are frequently made by comparison of the observed lengths with that predicted from Pauling's covalent radii (1.74 Å)<sup>[17]</sup> or with the bond length in sulfamic acid (1.772(1) Å).<sup>[15b]</sup> In the solid state, the latter has the zwitterionic structure <sup>+</sup>NH<sub>3</sub>–SO<sub>3</sub><sup>−</sup>, which possesses a formal S–N single bond. The observed S–N bond length in SO<sub>3</sub>NF<sub>2</sub><sup>−</sup> (1.772(4) Å) is similar to that in sulfamic acid and larger than the typical values found in the SO<sub>3</sub>NH<sub>2</sub><sup>−</sup> ion (1.64 Å). The S–O bond lengths (1.428(8) Å and 1.424(3) Å) are shorter than the values reported for sulfamates (1.44–1.45 Å)<sup>[16]</sup> and sulfamic acid (1.438(1) Å, 1.436(1) Å, and 1.435(1) Å).<sup>[15b]</sup> The average O–S–O and O–S–N angles (115.9° and 101.8°, respectively) are in good agreement with those reported for sulfamic acid.

The marked differences in the S–N and S–O bond lengths between SO<sub>3</sub>NF<sub>2</sub><sup>−</sup> and SO<sub>3</sub>NH<sub>2</sub><sup>−</sup> can be readily reconciled by the large difference between the electronegativity values of hydrogen and fluorine. The strongly electron withdrawing fluorine atoms pull some of the negative charge away from the oxygen atoms, thereby increasing the S=O double-bond character and shortening the S–O bonds. By contrast, the electron-donating hydrogen atoms increase the electron density on the nitrogen atom, which passes it on to the S=O bonds. This results in partial N=S double bond character and increased negative charge on the oxygen atoms with concomitant lengthening of the S–O bonds.

When only one equivalent of fluorine was used in the fluorination reaction of SO<sub>3</sub>NH<sub>2</sub><sup>−</sup>, the SO<sub>3</sub>NHF<sup>−</sup> ion was formed [Eq. (4)].



Colorless and somewhat moisture sensitive NaSO<sub>3</sub>NHF was obtained when the acidic reaction mixture was evaporated in a vacuum at 0°C. The compound was identified and characterized by its vibrational spectra and by its crystal structure as the salt [PPh<sub>4</sub>][SO<sub>3</sub>NHF]. The Raman and IR spectra of NaSO<sub>3</sub>NHF are shown in Supporting Information, and the observed frequencies and intensities are listed in Table 2 and were assigned by comparison with those calculated at the MP2<sup>[11]</sup> level of theory with the 6-311 + G(d) basis set.


**Table 2:** Comparison of observed and unscaled calculated MP2/6-311 + G(d) vibrational frequencies (cm<sup>−1</sup>) and intensities for SO<sub>3</sub>NHF<sup>−</sup> in point group C<sub>1</sub>.

Mode	Approx. mode description	Observed <sup>[a],[d]</sup>		Calculated <sup>[b]</sup>
		IR	Raman	(IR) [Raman]
$\nu_1$	$\nu$ NH	3100 s, br	3000 [1.3]	3481 (2.4) [69]
$\nu_2$	$\delta_{\text{sciss}}$ HNF	1522 w		1410 (30) [2.8]
$\nu_3$	$\nu_{\text{as}}$ SO <sub>3</sub>	1242 vs	1246 [1.3]	1270 (368) [7.3]
$\nu_4$	$\nu_{\text{as}}$ SO <sub>3</sub>	1213 vs	1225 [1.2]	1240 (401) [7.1]
$\nu_5$	$\nu_{\text{s}}$ SO <sub>3</sub> / $\delta_{\text{wag}}$ NH	1064 s	1065 [10.0]	1041 (82) [27]
$\nu_6$	$\nu_{\text{s}}$ SO <sub>3</sub> / $\delta_{\text{wag}}$ NH	1021 m	1014 [5.9]	1033 (45) [18]
$\nu_7$	$\nu$ NF	863 m	869 [4.3]	923 (38) [14]
$\nu_8$	$\nu$ SN	602 s	602 [5.0]	699 (161) [16]
$\nu_9$	$\delta_{\text{as}}$ SO <sub>3</sub>	590 s		584 (66) [4.5]
$\nu_{10}$	$\delta_{\text{as}}$ SO <sub>3</sub>	497 w	466 [2.6]	520 (26) [3.4]
$\nu_{11}$	$\delta_{\text{sym}}$ SO <sub>3</sub>	477 w	445 [3.0]	509 (26) [5.9]
$\nu_{12}$	$\delta_{\text{rock}}$ SO <sub>3</sub>	438 w	424 [3.7]	371 (0.1) [4.7]
$\nu_{13}$	$\delta_{\text{rock}}$ SO <sub>3</sub>		411 [3.0]	343 (8.1) [2.6]
$\nu_{14}$	$\delta_{\text{wag}}$ NF		[c]	215 (1.2) [1.1]
$\nu_{15}$	$\tau$ S–N		[c]	118 (1.9) [0.4]

[a] As Na<sup>+</sup> salt. Relative Raman intensities given in brackets. [b] IR intensities given in km mol<sup>−1</sup> (in parentheses) and Raman intensities given in Å<sup>4</sup> amu<sup>−1</sup> (in brackets). [c] Not observed, IR spectrum recorded only between 4000 and 400 cm<sup>−1</sup>. [d] In addition to the listed bands, IR bands at 1313 m, 1168 s, 1082 s, 918 m, 897 m, 773 m, 731 s, 617 m and Raman bands at 1168 [1.0], 1153 [1.0], 913 [1.2], 666 [1.0] cm<sup>−1</sup> were observed but were not assigned.

The [PPh<sub>4</sub>][SO<sub>3</sub>NHF] salt was obtained by metathesis in the same manner as shown in Equation (3). It crystallizes in the triclinic space group *P* $\bar{1}$ . The X-ray structure analysis reveals the presence of two nonequivalent [PPh<sub>4</sub>]<sup>+</sup> and [SO<sub>3</sub>NHF]<sup>−</sup> units (packing diagrams are shown in the Supporting Information). One SO<sub>3</sub>NHF<sup>−</sup> ion could be refined well, but the SO<sub>3</sub> group of the second ion is disordered. This results in a relatively high *R* factor of 8.16% for the refined structure. The closest F...H and O...H contacts between neighboring cations and anions are 2.566 Å and 2.373 Å, respectively. The ordered SO<sub>3</sub>NHF<sup>−</sup> ion is depicted in Figure 2. The observed S–N bond length in SO<sub>3</sub>NHF<sup>−</sup> (1.694(5) Å) lies between those observed for SO<sub>3</sub>NF<sub>2</sub><sup>−</sup> (1.772(4) Å) and SO<sub>3</sub>NH<sub>2</sub><sup>−</sup> (1.64 Å). This situation is in accord with the expectation that the electronic effect of an -NHF group is intermediate between those of an -NF<sub>2</sub> and an -NH<sub>2</sub> group.

The presence of one hydrogen atom and one fluorine atom attached to the same nitrogen atom was further established by proton and fluorine NMR spectroscopy. The



**Figure 2.** ORTEP drawing of the anion of  $[\text{PPh}_4][\text{SO}_3\text{NHF}]$ . Thermal ellipsoids are shown at the 50% probability level. Selected bond lengths [Å] and angles [°]: S–O1 1.433(4), S–O2 1.457(4), S–O3 1.447(3), S–N 1.694(5), F1–N 1.474(5), O1–S–O2 114.4(2); O1–S–O3 115.3(2), O2–S–O3 114.1(2), O1–S–N 108.5(2), O2–S–N 104.3(2), O3–S–N 98.0(2), F–N–S 106.4(3).

proton and fluorine spectra showed doublets at  $\delta = 8.69$  and  $-103.0$  ppm, respectively, with the same  $^2J(^1\text{H}-^{19}\text{F})$  coupling constant (49 Hz).

The utility of  $\text{NaSO}_3\text{NF}_2$  as an alternative source of difluoroamine in organic difluoroamination reactions was demonstrated by the transformation of carbonyl into *gem*-bis(difluoroamino) groups. Qualitative reactions of acetone, cyclohexanone, 3-pentanone, and 1-acetylpiperidin-4-one with the reagent in approximately 100%  $\text{H}_2\text{SO}_4$  and 30% oleum showed formation of the expected *gem*-bis(difluoroamino)alkane by  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{19}\text{F}$  NMR spectroscopy. In the case of 1-acetylpiperidin-4-one, only the oxygen atom of the ring-carbonyl group was replaced, whereas the carbonyl oxygen atom of the acetyl group did not react. A quantitative comparison was conducted by using 1,3-dibromoacetone as a model ketone. 1,3-Dibromo-2,2-bis(difluoroamino)propane was formed in 38% yield by using  $\text{NaSO}_3\text{NF}_2$  in anhydrous  $\text{H}_2\text{SO}_4$  containing approximately 1%  $\text{SO}_3$ . This compares with a yield of 37% for conventional difluoroamination ( $\text{HNF}_2$  in oleum) of 1,3-dibromoacetone.

## Experimental Section

**Materials and Apparatus (USC):** All reactions were carried out in teflon-FEP ampules closed by stainless steel valves or pyrex glass vessels closed by grease-free Kontes glass-teflon valves. Volatile materials were handled in stainless steel/teflon-FEP<sup>[18]</sup> or pyrex glass vacuum lines. Nonvolatile solids were handled in the dry argon atmosphere of a glove box. Infrared spectra were recorded in the range 4000–400  $\text{cm}^{-1}$  on a Midac, M Series, FT-IR spectrometer using KBr pellets. The pellets were prepared inside the glove box using an Econo mini-press (Barnes Engineering Co.) and transferred in a closed container to the spectrometer before placing them quickly into the sample compartment, which was purged with dry nitrogen to minimize exposure to atmospheric moisture and potential hydrolysis of the sample. Raman spectra were recorded on a Bruker Equinox 55

FT-RA spectrometer using a Nd-YAG laser at 1064 nm with power levels of 100–200 mW and pyrex melting-point capillary tubes as sample containers. NMR spectra were recorded unlocked on Bruker AMX 500 and Varian Mercury/VX 400 NMR spectrometers at room temperature. The  $^{19}\text{F}$  and  $^{15}\text{N}$  NMR spectra were referenced to external samples of  $\text{CFCl}_3$  and nitromethane, respectively.

The starting materials  $\text{H}_2\text{NSO}_3\text{H}$ ,  $\text{H}_2\text{SO}_4$ ,  $[\text{P}(\text{C}_6\text{H}_5)_4]\text{Cl}$ ,  $[\text{PNP}]\text{Cl}$ ,  $\text{NaOH}$  (all Aldrich),  $^{15}\text{N}_2$  urea (MSD Isotopes), and fluorine (Air Products and Chemicals Inc.) were used without further purification. Solvents were dried by standard methods and freshly distilled prior to use.  $\text{NaSO}_3\text{NH}_2$  was obtained by neutralization of  $\text{HSO}_3\text{NH}_2$  with  $\text{NaOH}$ .  $^{15}\text{N}[\text{HSO}_3\text{NH}_2]$  was prepared from  $^{15}\text{N}_2$  urea and  $\text{H}_2\text{SO}_4$  according to a modified literature method.<sup>[19]</sup>

**$\text{NaSO}_3\text{NHF}$  (USC):** A solution of  $\text{NaSO}_3\text{NH}_2$  (0.380 g, 3.19 mmol) in water (4 mL) was placed into a teflon-FEP ampule equipped with a teflon-coated magnetic stirring bar and a teflon gas-inlet tube. After cooling to 0°C, fluorine (3.19 mmol; 10% v/v in nitrogen) was introduced at a rate of 110  $\text{mL min}^{-1}$ . The reaction mixture was pumped to dryness at 0°C, leaving behind a colorless solid (0.391 g; weight calcd for 3.19 mmol  $\text{NaSO}_3\text{NHF} = 0.437$  g).

**$[\text{PPh}_4][\text{SO}_3\text{NHF}]$  (USC):** A solution of  $\text{NaSO}_3\text{NH}_2$  (0.695 g, 5.00 mmol) in water (10 mL) was placed into a teflon-FEP ampule equipped with a magnetic stirring bar and a teflon gas-inlet tube. After cooling to 0°C, fluorine (5.00 mmol; 10% v/v in nitrogen) was introduced at a rate of 110  $\text{mL min}^{-1}$ . The reaction mixture was neutralized by adding cold solid  $\text{NaHCO}_3$  and filtered through a cold porcelain frit. A cold solution of  $\text{PPh}_4\text{Cl}$  (5.00 mmol) in water (50 mL) was added to the clear filtrate. The resulting milky solution was extracted three times with 50 mL of cold  $\text{CH}_2\text{Cl}_2$ . The combined organic phases were dried over  $\text{MgSO}_4$  and pumped to dryness at ambient temperature, leaving behind a colorless solid (2.039 g; weight calcd for 5.00 mmol  $\text{PPh}_4\text{SO}_3\text{NHF} = 2.267$  g). Crystals were grown from a solution in  $\text{CH}_2\text{Cl}_2$  by slow evaporation of the solvent with a stream of dry nitrogen.

**$\text{NaSO}_3\text{NF}_2$  (USC):** A solution of  $\text{NaSO}_3\text{NH}_2$  (0.236 g, 1.98 mmol) in water (7 mL) was placed into a teflon-FEP ampule equipped with a magnetic stirrer and a teflon gas-inlet tube. After cooling to 0°C, fluorine (10% v/v in nitrogen) was introduced at a rate of 110  $\text{mL min}^{-1}$ . After 45 min, the fluorination was discontinued, and the reaction mixture pumped to dryness at 0°C, leaving behind a colorless solid (0.288 g; weight calcd for 1.98 mmol  $\text{NaSO}_3\text{NF}_2 = 0.307$  g).

**$\text{NaSO}_3\text{NF}_2$  (TPL):** Sulfamic acid (16.42 g, 0.169 mol) was dissolved in  $\text{H}_2\text{O}$  (30 mL), and  $\text{NaOH}$  (8.1 g, 0.20 mol) was added. Fluorine (10% v/v in nitrogen) was bubbled through the solution at 0°C. The progress of the fluorination of sulfamate was monitored by  $^{19}\text{F}$  NMR spectroscopy of reaction aliquots to observe the conversion of monofluorosulfamate ( $\delta = -102.1$  ppm vs. external  $\text{CFCl}_3$  in  $[\text{D}_6]\text{acetone}$ ) into difluorosulfamate ( $\delta = 36.6$  ppm). After 1.5 h, the fluorination was complete, and the cloudy mixture was neutralized with concentrated aqueous  $\text{NaOH}$ . The  $\text{NaF}$  precipitate was removed by filtration. On the basis of its  $^{19}\text{F}$  NMR spectrum, the filtrate appeared to be a 96:4 mixture of  $\text{NaSO}_3\text{NF}_2$  and  $\text{NaF}$ . The viscous filtrate was quickly dried at room temperature in a vacuum desiccator over excess  $\text{P}_2\text{O}_5$ . After 24 h,  $^{19}\text{F}$  NMR spectroscopy of the dried solid showed the same  $\text{SO}_3\text{NF}_2^-/\text{F}^-$  ratio as the initial filtrate.

**$[\text{A}][\text{SO}_3\text{NF}_2]$  (A = PNP,  $\text{PPh}_4$ ,  $\text{AsPh}_4$ ) (USC):** A solution of  $\text{NaSO}_3\text{NH}_2$  (0.357 g, 3.00 mmol) in water (10 mL) was placed into a teflon-FEP ampule equipped with a magnetic stirring bar and a teflon gas-inlet tube. After cooling to 0°C, fluorine (10% v/v in nitrogen) was introduced at a rate of 110  $\text{mL min}^{-1}$ . After 60 min, the fluorination was discontinued, and the reaction mixture was neutralized by adding cold solid  $\text{NaHCO}_3$ . The reaction mixture was filtered through a cold porcelain frit, and a cold aqueous solution of 3.00 mmol  $\text{A}^+\text{Cl}^-$  was added to the clear filtrate. The resulting milky solution was extracted three times with 10 mL of cold  $\text{CH}_2\text{Cl}_2$ . The combined organic phases were dried over  $\text{MgSO}_4$  and pumped to



dryness at ambient temperature, leaving behind colorless solids ([PPh<sub>4</sub>][SO<sub>3</sub>NF<sub>2</sub>]: 1.283 g, calcd weight for 3.00 mmol = 1.414 g; [AsPh<sub>4</sub>][SO<sub>3</sub>NF<sub>2</sub>]: 1.466 g, calcd weight for 3.00 mmol = 1.543 g; [PNP][SO<sub>3</sub>NF<sub>2</sub>]: 2.071 g, calcd weight for 3.00 mmol = 2.011 g). Crystals were grown from a solution in CH<sub>2</sub>Cl<sub>2</sub> by slow evaporation of the solvent with a stream of dry nitrogen.

2,2-Bis(difluoroamino)propane and 1-acetyl-4,4-bis(difluoroamino)piperidine (USC): NaSO<sub>3</sub>NF<sub>2</sub> (2.5 equiv.) was added in increments at 4 °C to a vigorously stirred mixture of CDCl<sub>3</sub> (1.5 mL), 30 % oleum (1.5 mL), and the ketone starting material (1.0 mmol). Immediate gas evolution was observed. After 15 min, the upper organic phase was analyzed by <sup>19</sup>F, <sup>1</sup>H, and <sup>13</sup>C NMR spectroscopy, showing the formation of the desired *gem*-bis(difluoroamines).

1,3-Dibromo-2,2-bis(difluoroamino)propane (TPL): 1,3-Dibromoacetone (24.72 g, 0.114 mol) was added to a mixture of sodium difluorosulfamate (61 g, ≤ 0.39 mol) in sulfuric acid (310 mL) containing approximately 1 % SO<sub>3</sub> (made from 30 % oleum plus concd H<sub>2</sub>SO<sub>4</sub>) plus 100 mL dichloromethane. The reaction mixture was stirred for 4 days at 0 °C, quenched in water, neutralized, and extracted with dichloromethane. Removal of the solvent produced 13.2 g of solute, the major constituent of which was 1,3-dibromo-2,2-bis(difluoroamino)propane (38 % crude yield). Three short-path distillations (room temperature at 1–3 Torr) produced material that was > 95 % pure (elemental analysis and <sup>1</sup>H NMR) with residual 1,3-dibromoacetone as the main impurity. <sup>1</sup>H NMR spectroscopy (200 MHz, CDCl<sub>3</sub>, TMS): δ = 3.98 ppm (quint., 1.25 Hz). <sup>13</sup>C{<sup>1</sup>H} NMR spectroscopy (CDCl<sub>3</sub>): δ = 23.75 (quint., 5.5 Hz), 99 ppm (m). <sup>19</sup>F NMR spectroscopy (CDCl<sub>3</sub>): δ = 29.9 ppm (s). <sup>1</sup>H NMR spectroscopy (C<sub>6</sub>D<sub>6</sub>): δ = 3.24 ppm (quint., 1.27 Hz). <sup>13</sup>C NMR spectroscopy (C<sub>6</sub>D<sub>6</sub>): δ = 23.69 (quint., 5.7 Hz), 95.37 ppm (quint., 6.4 Hz). <sup>19</sup>F NMR spectroscopy (C<sub>6</sub>D<sub>6</sub>): δ = 30.26 ppm (s). Elemental analysis calcd for (0.953)C<sub>3</sub>H<sub>4</sub>F<sub>4</sub>N<sub>2</sub>Br<sub>2</sub> + (0.047)C<sub>3</sub>H<sub>4</sub>Br<sub>2</sub>O: C 12.08, H 1.35, N 8.78, F 23.83; found: C 12.12, H 1.30, N 8.93, F 23.53 %.

Theoretical Methods: The molecular structures and harmonic vibrational frequencies were calculated by using second-order many-body perturbation theory<sup>[14]</sup> (denoted as MP2, but also known as MBPT(2)) and a 6-311+G(d) basis set. Hessians (energy second derivatives) were calculated for the final equilibrium structures to determine if they were minima (positive definite hessian) or *n*th-order transition states (“*n*” negative eigenvalues). All calculations were performed by using the electronic structure code GAMESS.<sup>[20]</sup>

X-ray structure determination: The collection, reduction, and processing of X-ray data was described previously.<sup>[21]</sup> Crystal data for C<sub>37</sub>H<sub>32</sub>Cl<sub>2</sub>F<sub>2</sub>N<sub>2</sub>O<sub>3</sub>P<sub>2</sub>S: *M*<sub>r</sub> = 755.55, triclinic, space group *P*1̄, *a* = 11.8650(12), *b* = 11.9167(12), *c* = 14.5180(15) Å, α = 66.755(2), β = 68.964(2), γ = 71.034(2)°, *V* = 1719.7(3) Å<sup>3</sup>, *F*(000) = 780, ρ<sub>calcd</sub> (*Z* = 2) = 1.459 g cm<sup>−3</sup>, μ = 0.395 mm<sup>−1</sup>, approximate crystal dimensions 0.29 × 0.21 × 0.05 mm<sup>3</sup>, θ range = 1.58 to 27.49°, index ranges −15 ≤ *h* ≤ 12, −15 ≤ *k* ≤ 12, −18 ≤ *l* ≤ 18, MoKα (λ = 0.71073 Å), *T* = 133(2) K, 10657 measured reflections, of which 7433 (*R*<sub>int</sub> = 0.0487) were unique, data to parameters ratio = 16.9:1, final *R* indices [*I* > 2σ(*I*): *R*1 = 0.0731, *wR*2 = 0.1949, *R* indices (all data): *R*1 = 0.1087, *wR*2 = 0.2102, GOF on *F*<sup>2</sup> = 1.012.

Crystal data for C<sub>37</sub>H<sub>32</sub>Cl<sub>2</sub>F<sub>2</sub>N<sub>2</sub>O<sub>3</sub>P<sub>2</sub>S: *M*<sub>r</sub> = 755.55, triclinic, space group *P*1̄, *a* = 11.8650(12), *b* = 11.9167(12), *c* = 14.5180(15) Å, α = 66.755(2), β = 68.964(2), γ = 71.034(2)°, *V* = 1719.7(3) Å<sup>3</sup>, *F*(000) = 780, ρ<sub>calcd</sub> (*Z* = 2) = 1.459 g cm<sup>−3</sup>, μ = 0.395 mm<sup>−1</sup>, approximate crystal dimensions 0.29 × 0.21 × 0.05 mm<sup>3</sup>, θ range = 1.58 to 27.49°, index ranges −15 ≤ *h* ≤ 12, −15 ≤ *k* ≤ 12, −18 ≤ *l* ≤ 18, MoKα (λ = 0.71073 Å), *T* = 133(2) K, 10657 measured reflections, of which 7433 (*R*<sub>int</sub> = 0.0487) were unique, data to parameters ratio = 16.9:1, final *R* indices [*I* > 2σ(*I*): *R*1 = 0.0731, *wR*2 = 0.1949, *R* indices (all data): *R*1 = 0.1087, *wR*2 = 0.2102, GOF on *F*<sup>2</sup> = 1.012.

CCDC-274108 and -274109 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge

from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

Received: March 15, 2006

Published online: July 6, 2006

**Keywords:** difluoroamines · fluorine · structure elucidation · sulfamates

- a) R. D. Chapman, C. B. Kreutzberger, M. F. Welker, *JANNAF Novel Ingredients for Liquid and Solid Propellants Specialist Session*, Pasadena, CA, 4 April, 1995; b) R. D. Chapman, R. Y. Yee, R. D. Gilardi, A. A. Pinkerton, *15th Annual Working Group Institute on Synthesis of High Energy Density Materials*, Picatinny Arsenal, NJ, 12–13 June 1996; available from the National Technical Information Service (<http://www.ntis.gov/help/ordermethods.asp>).
- a) R. D. Chapman, M. F. Welker, C. B. Kreutzberger, *J. Org. Chem.* **1998**, 63, 1566; b) R. D. Chapman, R. D. Gilardi, M. F. Welker, C. B. Kreutzberger, *J. Org. Chem.* **1999**, 64, 960.
- T. Axenrod, X.-P. Guan, J. Sun, L. Qi, R. D. Chapman, R. D. Gilardi, *Tetrahedron Lett.* **2001**, 42, 2621.
- a) K. Baum, *J. Am. Chem. Soc.* **1968**, 90, 7083; b) W. H. Graham, J. P. Freeman, *J. Org. Chem.* **1969**, 34, 2589; c) A. V. Fokin, Y. M. Kosyrev, V. A. Makarov, N. P. Novoselov, *Dokl. Akad. Nauk SSSR* **1969**, 186, 112.
- E. W. Lawless, I. C. Smith, *Inorganic High Energy Oxidizers (Syntheses, Structure and Properties)*, Marcel Dekker, New York, NY, **1968**, p. 69.
- J. K. Ruff, *Chem. Rev.* **1967**, 67, 665, and references therein.
- J. P. Freeman, A. Kennedy, C. B. Colburn, *J. Am. Chem. Soc.* **1960**, 82, 5304.
- a) V. Grakauskas, K. Baum, *J. Am. Chem. Soc.* **1970**, 92, 2096; b) R. E. Banks, R. N. Haszeldine, J. P. Lahu, *J. Chem. Soc. C* **1966**, 1514.
- a) V. Grakauskas, K. Baum, *J. Am. Chem. Soc.* **1969**, 91, 1679; b) J. M. J. Lambert, G. M. Alain (Société Nationale des Poudres et Explosives), DE 2320119, **1973**; c) D. L. Klopotek, G. B. Hobrock, *Inorg. Chem.* **1967**, 6, 1751.
- a) R. C. Petry, J. P. Freeman, *J. Am. Chem. Soc.* **1961**, 83, 3912; b) W. H. Graham, C. O. Parker, *J. Org. Chem.* **1963**, 28, 850; c) G. K. S. Prakash, M. Etzkorn, G. A. Olah, K. O. Christe, S. Schneider, A. Vij, *Chem. Commun.* **2002**, 1712.
- J. N. Keith, R. J. Douthart, W. K. Sumida, I. J. Solomon in *Advanced Propellant Chemistry* (Ed.: R. T. Holzmann), American Chemical Society, Washington, **1966**, pp. 141–147.
- R. D. Shoults, *Development of a Practical Process for the Preparation of gem Difluoroamino Compounds. The Difluoro-sulfamic Acid Reagent*, Rohm and Haas Co. Report S-129, **1967**, NTIS Accession Number AD0380719, available from the National Technical Information Service (<http://www.ntis.gov/help/ordermethods.asp>).
- R. A. Wiesboeck, J. K. Ruff, *Inorg. Chem.* **1965**, 4, 123.
- a) C. Möller, M. S. Plesset, *Phys. Rev.* **1934**, 46, 618; b) J. A. Pople, J. S. Binkley, R. Seeger, *Int. J. Quantum Chem. Symp.* **1976**, 10, 1; c) M. J. Frisch, M. Head-Gordon, J. A. Pople, *Chem. Phys. Lett.* **1990**, 166, 275; d) J. Bartlett, D. M. Silver, *Int. J. Quantum Chem. Symp.* **1975**, 9, 1927.
- a) F. A. Kanda, A. J. King, *J. Am. Chem. Soc.* **1951**, 73, 2315; b) A. F. Cameron, F. D. Duncanson, *Acta Crystallogr. Sect. B* **1976**, 32, 1563.
- a) V. K. Wadahawan, V. M. Padmanabhan, *Acta Crystallogr. Sect. B* **1972**, 28, 1903; b) R. Manickavachagam, R. K. Rajaram, *Z. Kristallogr.* **1984**, 168, 179; c) A. Meinhart, E. Haussühl, L. Bohatý, E. Tillmanns, *Z. Kristallogr.* **2001**, 216, 513; d) J. Schreuer, *Z. Kristallogr. New Cryst. Struct.* **1999**, 214, 305; e) J.

- Schreuer, Z. *Kristallogr. New Cryst. Struct.* **1999**, 214, 306; f) J. Schreuer, Z. *Kristallogr. New Cryst. Struct.* **1999**, 214, 307; g) J. Schreuer, Z. *Kristallogr. New Cryst. Struct.* **1999**, 214, 309; h) M. S. Wickleder, Z. *Anorg. Allg. Chem.* **1999**, 625, 1794.
- [17] L. Pauling, *The nature of the chemical bond*, 3rd ed., Cornell University Press, Ithaca, NY, **1960**, p. 310.
- [18] K. O. Christe, W. W. Wilson, C. J. Schack, R. D. Wilson, *Inorg. Synth.* **1986**, 24, 39.
- [19] P. Baumgarten, *Ber. Dtsch. Chem. Ges. B* **1936**, 69, 1929.
- [20] M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, *J. Comput. Chem.* **1993**, 14, 1347.
- [21] K. O. Christe, M. Gerken, R. Haiges, S. Schneider, T. Schroer, I. Tsyba, R. Bau, *Inorg. Chem.* **2003**, 42, 416.